

	pT2	pT3	pT4	p-value
<b>TILs</b>				
None/low	6 (85.7)	141 (76.2)	18 (81.8)	0.661
Intermediate	1 (14.3)	30 (16.2)	4 (18.2)	
High	0 (0.0)	14 (7.6)	0 (0.0)	
<b>Mucus</b>				
≤50%	7 (100)	165 (89.2)	17 (77.3)	0.160
>50%	0 (0.0)	20 (10.8)	5 (22.7)	
<b>Mucus</b>				
<10%	4 (57.1)	131 (70.8)	13 (59.1)	0.416
≥10%	3 (42.9)	54 (29.2)	9 (40.9)	
<b>TSR</b>				
Stroma-low	7 (100)	125 (66.1)	13 (59.1)	0.130
Stroma-high	0 (0.0)	64 (33.9)	9 (40.9)	
<b>Tumor Budding</b>				
Low (<5)	7 (100)	150 (79.4)	19 (86.4)	0.439
Intermediate (5-9)	0 (0.0)	28 (14.8)	1 (4.5)	
High (≥10)	0 (0.0)	11 (5.8)	2 (9.1)	

**Table S3.** Correlation between histopathologic features and pathologic T-stage. TILs = tumor infiltrating lymphocytes, TSR = tumor-stroma ratio. P-values are derived from an overall comparison between pathologic T-stages.