


Deconstructing the mouse olfactory percept through an ethological atlas

Graphical abstract



Authors

Diogo Manoel, Melanie Makhoulouf, Charles J. Arayata, ..., Joel D. Mainland, Richard C. Gerkin, Luis R. Saraiva

Correspondence

rgerkin@asu.edu (R.C.G.), saraivalmr@gmail.com (L.R.S.)

In brief

Manoel et al. score 18 behaviors in 525 mice across 98 odorant conditions to generate a murine olfactory ethological atlas, which they use to deconstruct mouse olfactory behavior, predict it from odorant structure, evaluate its major factors, and compare it to human perception.

Highlights

- We produced a novel atlas of mouse odor-guided behavior for 73 molecules
- Odorant structure and mouse behavior are mutually predictive in novel mice
- Odor valence explains most behavioral variance and covariance with human perception
- Concentration qualitatively and quantitatively impacts mouse olfactory behavior



Article

Deconstructing the mouse olfactory percept through an ethological atlas

Diogo Manoel,¹ Melanie Makhoul,¹ Charles J. Arayata,² Abbirami Sathappan,¹ Sahar Da'as,¹ Doua Abdelrahman,¹ Senthil Selvaraj,¹ Reem Hasnah,¹ Joel D. Mainland,^{2,3} Richard C. Gerkin,^{4,*} and Luis R. Saraiva^{1,2,5,6,7,*}

¹Sidra Medicine, PO Box 26999, Doha, Qatar

²Monell Chemical Senses Center, 3500 Market Street, Philadelphia, PA 19104, USA

³Department of Neuroscience, University of Pennsylvania, Philadelphia, PA 19104, USA

⁴School of Life Sciences, Arizona State University, Tempe, AZ 85281, USA

⁵College of Health and Life Sciences, Hamad Bin Khalifa University, Doha, Qatar

⁶Twitter: @saraivalab

⁷Lead contact

*Correspondence: rgerkin@asu.edu (R.C.G.), saraivalmr@gmail.com (L.R.S.)

<https://doi.org/10.1016/j.cub.2021.04.020>

SUMMARY

Odor perception in non-humans is poorly understood. Here, we generated the most comprehensive mouse olfactory ethological atlas to date, consisting of behavioral responses to a diverse panel of 73 odorants, including 12 at multiple concentrations. These data revealed that mouse behavior is incredibly diverse and changes in response to odorant identity and concentration. Using only behavioral responses observed in other mice, we could predict which of two odorants was presented to a held-out mouse 82% of the time. Considering all 73 possible odorants, we could uniquely identify the target odorant from behavior on the first try 20% of the time and 46% within five attempts. Although mouse behavior is difficult to predict from human perception, they share three fundamental properties: first, odor valence parameters explained the highest variance of olfactory perception. Second, physicochemical properties of odorants can be used to predict the olfactory percept. Third, odorant concentration quantitatively and qualitatively impacts olfactory perception. These results increase our understanding of mouse olfactory behavior and how it compares to human odor perception and provide a template for future comparative studies of olfactory percepts among species.

INTRODUCTION

How sensory cues translate into perceptual objects or complex behaviors remains a major unanswered question in neuroscience. Odor transduction in the nose leads to odor perception and to changes in behavior or physiology (e.g., aggression and feeding) that are key for survival and reproduction, making the olfactory system an attractive model to address this question.^{1,2}

Many studies have used mice to elucidate molecular, cellular, and neural processes underlying mammalian olfaction.¹ The proliferation of annotated genomes and high-throughput sequencing technologies have yielded new clues into the functional logic and the evolutionary dynamics of mammalian olfaction more broadly.^{3–5} However, our understanding of olfactory perception derives from large psychophysical datasets in humans combined with chemoinformatic, statistical, and machine-learning tools.^{6–11} These studies have yielded three key findings regarding human olfactory perception. First, the first principal component of human olfactory perception is highly associated ($r = 0.79$) with a single dimension—odor valence.^{6,7,12} Second, the human olfactory perceptual ratings for most odorous molecules can be predicted from chemical structure with surprising accuracy ($r = 0.3–0.7$), a value limited primarily

by rater reliability.⁸ Third, odorant concentration can qualitatively alter perceived odor intensity and character.^{9,13} Do these principles also apply to olfactory perception in non-humans?

Characterizing olfactory perception in an animal relies on accurately quantifying multiple behaviors in response to large numbers of odorants, ideally at various concentrations. This assumes that visible mouse behaviors either encode for mouse perception or at least report something informative about the meaning of the stimulus. For example, mice may exhibit differential behavioral responses depending on odor valence (attractive versus aversive), odor novelty, or implications of the presence of an odor for a broader behavioral strategy (e.g., exploration of surroundings). They may also use behavior to communicate odor information to conspecifics. Despite recent efforts,^{14–25} a systematic characterization of various mouse behaviors in response to a large panel of diverse odorants and several concentrations is still lacking. This prevents a systematic understanding of mouse olfactory behavior and how it relates to perception in humans and other species and limits our ability to study the neural computations underlying the transformation of odor stimuli at the nose to odor objects in the brain.

Here, we generated and investigated a mouse atlas of odor-guided behaviors in response to a diverse panel of odorants, at






Figure 1. An olfactory-ethological atlas and the primary axis of olfactory perception in mouse

(A–D) Videos (A) were scored for 18 behavioral parameters, grouped in three broad categories: (B) valence, (C) stress, and (D) exploration.

(E) A graphical display summarizing the combinatorial behavioral patterns for the 73 odors tested. Behaviors showing significant (one-way ANOVA; BKJ multiple comparisons correction; $n = 5–10$ per odorant) increases, decreases, and non-significant responses compared to H₂O are indicated in green, magenta, and gray squares, respectively.

(F) Correlation across odors between each of the 18 behaviors ($n = 3–7$ per odorant).

(G) Correlation across behaviors between each of the 73 odors and H₂O ($n = 3–7$ per odorant).

In both (F) and (G), odors and behaviors are ordered to illustrate clustering. Two major clusters of odors stand out.

(H) Principal-component analysis of the 18 behaviors for H₂O and the 73 tested odors ($n = 3–7$ per odorant). Circles are colored to indicate avoidance (magenta), neutral (gray), or approach (green) odors.

See also [Figure S1](#) and [Data S1](#).

different concentrations, to deconstruct olfactory behavior in mice and compare it to human perception.

RESULTS

The olfactory-ethological atlas

We recently generated a mouse behavioral video library and quantified the cumulative duration of olfactory investigation over a 3-min-long assay (3'dOI) to a panel of 73 odors at

85 mM and the odorless control water (H₂O).²⁵ These stimuli include 61 general odors, seven pheromones (IPT, IAA, BFA, 2HO, TMA, AST, and FAR), and five kairomones (TMT, PEA, 2PT, BZL, and QUI).^{2,26–30} Odors were classified as eliciting avoidance or approach if the 3'dOI was significantly lower or higher than H₂O, respectively.²⁵ Here, we enhance the utility of this video library by scoring 17 additional behavioral parameters from 410 mice exposed to 73 odors and H₂O ([STAR Methods](#); [Figures 1A–1D](#)). Each mouse was exposed to only

one odorant ($n = 5$ – 10 per odorant) and scored on 18 distinct behavioral parameters, chosen for discernibility, quantifiability, and reliability and as surrogate measures of odor “valence,” exploration, or stress.^{14,15,20,23,25,31–34}

From this new dataset totaling 7,606 individual data points (Figure S1A; Data S1), corresponding to individual-odorant-behavior triads, we calculated the across-individual average value for each of 1,314 odorant-behavior pairs: 470 (35.8%) are significantly different ($p < 0.05$; one-way ANOVA; BKY multiple-comparisons correction) from H₂O, with 293 decreases and 177 increases of a behavioral parameter (Figure 1E; Data S1). Of the 72 odorants eliciting at least one significant behavioral change, 57 (78%) exhibited unique patterns of significant increases and decreases among the 18 behaviors, indicating that odor-guided behavioral patterns are incredibly diverse.

Valence is the primary axis of olfactory perception in mouse

Pleasantness, a surrogate for odor valence, is the primary axis of human olfactory perception.^{7,12,35} However, whether this is conserved in mouse remains unknown. We examined the structure of the behavioral response matrix (behaviors \times odorants) to address this question. From this matrix, we computed two correlation matrices (between behaviors [Figure 1F] and between odorants [Figure 1G]) and ordered each to match a hierarchical clustering. We identified two self-similar groups (upper left and lower right patches in Figures 1F and 1G). Principal-component analysis (PCA) showed that principal component 1 (PC1) explains 30.4% of the data variance (Figures 1H and S1B). We observed only a weak relationship between molecular structure and PC1 or PC2 (Figure S1C), suggesting that odor valence is not a trivial consequence of molecular structure. Nevertheless, PC1 cleanly separates approached from avoided odorants, which we interpret as a valence axis (Figure 1H). This result is robust to alternative subsamples of behaviors in which valence, exploration, and stress have equal representation; valence-related behaviors always comprise the top 5 highest weighted behaviors in PC1.

Discriminability of odorants using behaviors

Odorants appear to lie in stereotypical locations in a behavioral space defined by PC1 and PC2 of the response matrix (Figure 1H). We next asked how well odorants can be distinguished using the original 18-dimensional space. We computed D' , a measure of discriminability between two signals, for all odorant pairs. Higher D' values indicate that behavior more easily discriminates between two odorants; a value of 1.0 occurs when the mean behavioral difference between odorants equals the behavioral variability across mice exposed to the same odorant; it is thus also a measure of effect size.

D' for each behavior (across odorant pairs) ranged between 0.6 and 1.0; by contrast, shuffling odorant labels between mice resulted in lower D' values (0.5–0.6; Figure 2A). Every single behavior exhibited higher D' in the original than in the shuffled data (Figure 2B).

Because the behavioral code may be combinatorial, we also computed a multivariate D' , which uses all behaviors simultaneously. If most behavioral variability is mouse specific and not odorant specific, D' would be much less than 1. However, we

found that multivariate D' —although heterogeneous across odorant pairs (Figure 2C)—was on average 4 \times larger for the real data (mean $D' = 0.8$) than for shuffled data (Figure 2D). Thus, this behavioral ensemble can represent an odorant-specific behavioral code. D' can alternatively be expressed in terms of the area under the receiver-operator characteristic (ROC) curve (AUC) quantifying a sensitivity versus specificity tradeoff; under a Gaussian assumption, a $D' = 0.8$ corresponds to an AUC = 0.66 for a typical pair of odorants.

D' can overestimate discriminability when the number of replicates (individuals) per stimulus is small (so D' for shuffled data remains >0). Instead, we can ask how accurately a predictive model can select the correct odorant from out-of-sample observations of behavior, i.e., a mouse the model has not observed. We trained a linear discriminant classifier on all odorants but withheld one mouse per odorant for cross-validation testing. We asked the classifier to perform two tasks: first, to predict the correct odorant (out of 74 possibilities) given a new observation of behavior and, second, to predict the correct odorant (out of two choices: the correct odorant and one other chosen at random). In the first task, most odorants could be predicted at above chance (1/74) levels (Figures 2E and 2F). Some could even be predicted correctly $>30\%$ of the time (IND, IBT, and AST). In the second task, 67/74 odorants could be identified from behavior at above chance ($1/2$) levels, and 32/74 could be identified $>95\%$ of the time in the same comparison (Figures 2G and 2H). If we gave the classifier additional “shots on goal” in the first task, allowing 5 shots made 46% of odorants identifiable from the ensemble (Figure 2I). Thus, odor-evoked behavior for most odorants was stereotypical enough to help identify which odorant was presented to a novel mouse.

Reconstructing a low-dimensional space of mouse olfactory behavior

We recorded 18 distinct behaviors, but many of these behaviors are correlated (Figure 1F). Thus, the underlying dimensionality of the olfactory behavioral space may be $\ll 18$. PCA indicated that 90% of the variance was explained by ten dimensions (Figure S1B). However, this is likely to be an overestimate for two reasons: first, at least some of this variance is noise, driven by within-odorant, across-mouse behavioral variability and, second, PCA does not produce a natural decomposition of data in many applications.³⁶ A particular concern here is that PCA might represent valence as a single dimension, although the underlying concepts “aversive” and “approach” could be distinct perceptual categories that just happen to produce behaviors of opposite sign. To overcome the first concern, we asked how many dimensions are required to optimally represent each odorant’s behavioral phenotype. To address the second, we used non-negative matrix factorization (NMF),³⁶ a decomposition technique known for producing compact, intuitive, parts-based representations in diverse domains, including olfactory perception.³⁷ Specifically, we computed an NMF decomposition of behavior and asked for what number of factors the intraclass correlation coefficient (ICC) was maximized when projecting data from novel mice onto these factors. Here, the ICC measures the fraction of behavioral variance explained by odorant identity. Theory suggests that a low-factor NMF decomposition might denoise behavioral data by identifying and discarding noisy,




Figure 2. Discriminability of odors using behaviors and out-of-sample prediction of odor identity

(A) D' , a measure of discriminability between two odors, is greater for the data than for a shuffling (across mice) of the data.

(B) Same data as in (A), except shown for each behavior (circle) versus its corresponding shuffle. Error bars (inside circles) represent SEM taken over all odorant pairs.

(C) Using all behaviors simultaneously, the multivariate measure D' is computed.

(D) Multivariate D' is $\sim 4\times$ larger for the real data than for shuffled data.

(E–H) A linear discriminant analysis classifier was trained on all odors, using all but one mouse for each odorant. Predictive performance was evaluated for the remaining mice (one odorant each).

(E) A histogram of the probability that the correct odorant (out of 74 possibilities) is identified from a new mouse's behavior. The dashed red line reflects chance performance.

(F) Mean performance for each odorant; higher values mean the odorant is easier to uniquely identify from behavior.

(G) Similar to (E) but for classification of the true odorant against a random alternative odorant. Chance is now 50%, as reflected by the dashed red line.

(H) Mean performance for each odorant in (G). 1.0 means that behavior was always sufficient to identify the odorant versus any specific alternative odorant.

(I) Number of shots (guesses) that the classifier needs to determine the correct odorant (out of 74 possibilities) from novel mouse behavior. This value (solid blue line) is shown for all 74 odorants, ranked from fewest to greatest number of shots required. The orange dashed line represents chance performance. See also Figures S2 and S3.

irrelevant dimensions, thus increasing ICC. However, with too few factors, odorant-associated structure could be lost, reducing ICC. The number of factors that maximizes ICC thus reflects the most efficient decomposition of olfactory behavior.

We found that ICC was greatest for a 2-factor space ($ICC = 0.57$; Figure 3A), substantially higher than for the original 18-factor space ($ICC = 0.34$) or a 1-factor space ($ICC = 0.35$). A 2-factor space was thus optimal for explaining mouse behavioral variance in terms of odorant identity. By contrast, shuffling odorant labels across mice produced a consistently low value ($ICC = 0.2$) for all choices of factor number. To interpret these two factors, we examined the factor weights for each behavior (Figure 3B). The highest weighted factors corresponded to valence-related behaviors (handling and catching [H/C], olfactory investigation [OI], and zone assays [Z1/Z3]), followed by the exploratory (rearing [REA], digging [DIG], and distance covered [DIT]) and stress-related behaviors (risk assessment [RAS] and escape [ESC]). The weights for each factor were mostly orthogonal (Figure S2A; $r = -0.24$; $p = 0.35$; Fisher Z transformation test), but the positions of odorants along each factor were anti-correlated (Figure S2B; $r = -0.79$; $p < 0.0001$), indicating that these two factors are still largely capturing a single behavioral category and its opposite (i.e., approach and aversion). This is consistent with PCA—the valence axis is primary—but goes a step further by separating out an attractive from an aversive factor and showing that these factors (but no others) are shared identifiably in cross-animal comparisons. This provides a parts-based understanding of the fundamental units of olfactory behavior in mice.

Mouse versus human

Comparative studies of odor valence in mice and humans are scarce and have yielded conflicting results.^{17,38} Indeed, whether mice and humans share a common olfactory perceptual space is




Figure 3. Optimal reconstruction of mouse behavior and the alignment between mouse and human behavioral spaces

(A and B) The optimal reconstruction of mouse behavior requires few dimensions.

(A) Non-negative matrix factorization (NMF) is used to learn a low-dimensional (k , the number of measured behaviors) representation of the behavioral data. The intra-class correlation coefficient (ICC), reflecting the behavioral agreement within (versus across) odorants, is shown as a function of the number of factors used. Lower numbers of factors effectively denoise the data. Results for the data are shown in blue; results for the data with shuffled odorant labels are shown in orange. Eighteen factors would reflect independent contributions of each behavior. ICC is maximized for a 2-factor representation of behavior.

(B) Contributions of behaviors to the resulting 2 factors.

(C) Alignment of mouse and human behavioral spaces. Canonical correlation analysis co-aligns mouse behavioral features and human-provided descriptors for the same odorants. Canonical dimensions were computed using all but one odorant, and the remaining odorant was used to evaluate the correlation (Pearson) between mouse behavior and human percepts. Error bars represent standard deviation across held-out odorants. p values were computed by comparing to shuffled data.

(D) Comparing mouse behaviors and human percepts. Correlation matrix heatmap comparing the 18 mouse behavioral parameters with 21 human odor descriptors for all 73 odorants tested here (odor descriptors for 22 odorants taken from Keller and Vosshall⁹ and 51 predicted using the DREAM model⁸). Rows and columns are sorted to maximize clustering in the heatmap, revealing ~2 major clusters of perception and behavior.

See also Figures S2 and S3 and Table S1.

still unknown. To answer this question, we first compared our mouse behavioral dataset to the human-rated intensity and pleasantness reported in a previous study.⁹ For the 23 overlapping odorants between both studies, we found no significant correlation among the mouse- and human-rated parameters (Figure S3A), consistent with our previous study.³⁸ We then extended this analysis to include 19 additional human semantic descriptors (e.g., fishy and sweet)⁹ but found only seven significant correlations (out of a possible 475) encompassing five mouse behaviors and four human semantic descriptors (Figure S3A).

To identify shared structure between mouse olfactory behavior and human olfactory perception, 23 odorants may be insufficient. To make use of a larger sample, we used the data and machine-learning algorithms from the DREAM Olfaction Prediction Challenge⁸ to obtain predictions of human-rated intensity, pleasantness, and 19 semantic descriptors for the remaining 51 odorants tested in our mouse experiments (STAR Methods). Rather than identify simple correlations, we asked whether mouse behavioral space and human perceptual space could be projected onto a common basis. If so, they might simply be two views of a common mammalian olfactory perceptual space. We used canonical correlation analysis (CCA) to obtain this basis using all but one odorant and then asked whether this basis could identify shared inter-species structure using the remaining (out-of-sample) odorant. Our results (Figure 3C) identified a single dimension for this hypothesized shared structure (Pearson's $r = 0.50$; $p = 0.02$; shuffle test), with additional dimensions failing to capture any additional shared structure. This means that a one-standard-deviation change in the optimal linear combination of mouse behaviors is associated with a $\frac{1}{2}$ standard deviation change in an optimal linear combination of human percepts for novel odorants. What human percepts comprise this shared dimension? The factor weights are shown in Table S1. No single human percept dominates this dimension, and individually, each of them is not statistically significant ($p > 0.1$). Similarly, all human percepts have a Pearson correlation $r < 0.5$ versus either of the first two principal components of mouse behavioral space in Figure 1 (Figure S3B) or the first two NMF factors obtained from the same space in Figure 3 (Figure S3C). The correlation between specific human percepts and mouse behaviors is shown in Figures 3D and S3A. We conclude that the shared dimension is multifactorial: it cannot be easily reduced to a single percept or behavior.

Predicting behavior from chemical structure

Prior studies identified relationships between hydrocarbon chain length or chemical functional group of aliphatic odorants and their induced neural responses or perceptual similarity.^{39–41} For the twelve aliphatic odorants in our dataset, we found no apparent association between chemical functional groups and valence (Figure 4A) but observed a strong rank correlation between hydrocarbon chain length and 3'dOI ($r_s = 0.795$; $p = 0.0034$; Figure 4B; Data S2). We observed similar significant relationships for three additional valence-related olfactory investigation parameters (1'dOI, 1'fOI, and 3'fOI), but not for other behavioral parameters. However, a broader understanding of odor perception requires a more complete description of the stimulus.^{7,42} Indeed, robust correlations have been identified




Figure 4. Predicting mouse olfactory-driven behaviors

(A) Twelve *n*-aliphatic odorants in our dataset include molecules with varying lengths and different functional groups: amyl acetate (AAC), hexanethiol (HXT), hexylamine (HXA), hexanol (HXO), heptanol (HPO), heptanal (HPN), heptylamine (HEP), octanoic acid (OCA), octanol (OCO), octanal (OCN), octylamine (OCT), and octanethiol (OTT). Circles are colored to indicate aversive (magenta), neutral (gray), or approached (green) odorants.

(B) Correlation plot between the numbers of carbon atoms and the total duration of olfactory investigation (3'dOI), for the 12 odorants displayed in (A) ($n = 5$ –9 mice per odorant). The R^2 value for the linear regression, Spearman's correlation coefficient (r_s), and associated p values are indicated in the top left corner of the graph.

(C) Heatmap depicting the r_s between 1,536 physicochemical descriptors and the 18 mouse behavioral parameters. The physicochemical descriptors are sorted in descending order of the r_s for 3'dOI.

(D) Performance of the support vector regression model on the 18 behavioral parameters. Blue bars are for models trained on all but one mouse and tested on that mouse. Orange bars are for models trained on all but one odorant and tested on novel mice for that odorant. Black borders around the bars indicate significant associations ($p < 0.01$).

See also Figure S3 and Data S2.

between large sets of physicochemical descriptors and multiple olfactory perceptual qualities in human, as well as simple mouse behavioral measures, like investigation time.^{8,9,17,43}

To test whether mouse behaviors are related to distinct physicochemical descriptors, we retrieved 4,870 physicochemical Dragon descriptors for all odorants. After removing descriptors with near-zero variance or missing data, we calculated the correlations between the remaining subset of 1,536 physicochemical descriptors and the 18 behavioral parameters for all 410 individual mice (Figure 4C; Data S2). Of the possible 27,648 interactions, 29.5% (or 8,141) resulted in significant ($p < 0.05$) interactions, with the durations of self-grooming (dSGR) and handling and catching (dH/C) eliciting the minimum (16) and maximum (968) significant interactions, respectively.

Physicochemical descriptors can be used to reverse-engineer perceptual descriptors of odorants in humans.^{8,43,44}

Can mouse olfactory-driven behaviors also be predicted by the chemical structure of odorants? To answer this question, we used an unsupervised machine-learning approach (STAR Methods) evaluated on either held-out mice or held-out odorants. 17/18 behaviors could be predicted above chance ($p < 0.01$) using physicochemical features on held-out mice, with predictive accuracy varying greatly from 1'dOI ($r = 0.59$; $p < 1e-40$) to dSGR ($r = 0.02$; $p = 0.31$). Because within-odorant behavior across mice is highly correlated, a stricter test is to predict behavior for all mice on a held-out odorant. There, 7/18 behaviors could be predicted ($p < 0.01$; or 9/18, $p < 0.05$) from physicochemical descriptors, with the four zone parameters (indicating approach versus avoidance) having the strongest predictability ($r = 0.38$ – 0.50 ; $p < 0.001$; Figure 4D).

Behavioral effects of odorant concentration

In humans, the quality, valence, and intensity of odorants can change with concentration.^{11,13,45} For example, humans perceive (R)-1-p-menthen-8-thiol as a pleasant grapefruit odor at low concentrations but as an unpleasant sulfur odor at high concentrations. In mouse, the valence^{25,31} and intensity⁴⁶ of odorants can also be concentration dependent, but whether concentration affects other odor-guided behaviors remains largely unexplored. To address this question, we scored 123 videos from mice exposed to a subset of 12 odorants at two additional descending concentrations (850 μ M and 8.5 μ M; Figure S4A; Data S3). At the highest concentration tested (85 mM), odorants have diverse valence, with some eliciting avoidance (IBT, IAA, and TMT), no change (PEA, 2HO, DMP, VAN, SKA, and AMB), or approach (IND, TMA, and PUT; Figure 5A). However, at lower concentrations, all odorants were neutral or elicited approach (Figure 5A).²⁵ Scoring the 17 other behaviors for each concentration yielded 3,474 individual data points (Data S3). Of the $12 \times 3 \times 18 = 648$ odorant-concentration-behavior pairwise comparisons, 117 (18.1%) significantly differed from H₂O (Figures 5A and S4A; Data S3). 3'dOI elicited the highest (23) and dSGR the lowest (1) number of significant behavioral changes (Figure 5A; Data S3), consistent with odor valence driving the largest fraction of behavioral variation. Of the 12 odorants tested, 10 elicited significant behavioral changes for all concentrations tested, although the remaining 2 (DMP and PEA) did so for only 2 out of 3 concentrations. Across all odorants, 85 mM elicited the highest number of combined significant behavioral changes (60), followed by 8.5 μ M (30) and 850 μ M (27). We identified 28/36 unique combinations of significant behavioral changes: 12 for 85 mM; 8 for 850 μ M; and 8 for 8.5 μ M (Figure 5A).

We next estimated the role of concentration and odorant-concentration interactions in generating behavioral responses. ANOVA showed a main effect of either concentration or an interaction between odorant and concentration for nearly all behavioral parameters ($p < 0.005$ for 16/18; $p < 10^{-8}$ for 13/18). The effect size for odorant identity ($\eta^2 = 0.27 \pm 0.05$) was slightly greater than for odorant concentration ($\eta^2 = 0.18 \pm 0.04$), but the interaction between odorant identity and concentration was stronger than either one alone ($\eta^2 = 0.43 \pm 0.04$).




Figure 5. The effect of odorant concentration on behavior

(A) A graphical display summarizing the behavioral codes for the 12 odorants tested at three different concentrations. Behaviors showing significant (one-way ANOVA; BKY multiple comparisons correction; $n = 5-9$ mice per odorant) increases, decreases, and non-significant responses compared to H2O are indicated in green, magenta, and gray squares, respectively.

(B) The behavioral profiles (radar plots and Z scores) for H2O (top center) and the 12 odorants tested at three concentrations (85 mM, 850 μM, and 8.5 μM) are shown. The behavioral profiles for the odorants classified at 85 mM as aversive, neutral, and approached are shown in different shades of magenta, gray or black, and green, respectively. The corresponding descending concentrations are depicted in lighter hues of the same color, irrespective of their valence at 850 μM and 8.5 μM.

(C) Hierarchical clustering analysis of the behavioral profiles for the 12 odorants tested at the three different concentrations supported the existence of 2 clusters. Bootstrap values⁴⁷ are shown only for strongly supported nodes, i.e., nodes displaying bootstrap values >70 (total 100 replications).

(D) Plot comparing the Euclidean distances calculated for all pairwise inter-concentration comparisons (restricted to different concentrations within each odorant), inter-odorant comparisons (restricted to different odorants at the same concentration), and subsets of inter-odorant comparisons for each concentration separately (85 mM, 850 μM, and 8.5 μM). Asterisks indicate significant differences (one-way ANOVA; BKY multiple comparisons correction); ns, non-significant; * $p < 0.05$; *** $p < 0.001$.

See also Figure S4 and Data S3.

Thus, even at low concentrations, odorants impact mouse behavior and produce diverse behavioral responses. But how does mouse olfactory behavior change with concentration and how is odor identity preserved across concentrations?

To answer these questions, we compared the behavioral profiles across odorants and concentrations. The highest concentration (85 mM) yielded the most distinguishable behavioral profiles across odorants, and some of these profiles were conserved for the same odorants at 850 μM and 8.5 μM (Figure 5B). As concentration decreased, the scores for some behaviors associated with stress (RAS and ESC) and negative odor valence (3'Z1) typically decreased, although the ones associated with positive odor valence (e.g., dH/C, 1'dOI, 3'dOI, and 3'Z3) increased.

Next, we performed a hierarchical clustering of odorants-concentrations pairs using the behavioral data. This analysis yielded two clusters (Figure 5C). Cluster 1 is composed mostly (7/8) of odorants presented at 85 mM (five neutral and three elicit avoidance), which elicit on average 7.00 ± 0.85 (SEM) significant behavioral changes. By contrast, cluster 2 includes H2O, eight neutral, and 20 approached odorants. These odorants elicit on average only 2.18 ± 0.28 (SEM) significant behavioral changes. Moreover, in cluster 2, five odorants (TMA, DMP, PUT, TMT, and VAN) produce behaviors which, although diverse across odorants, were quite similar across the two lowest

concentrations of the same odorant. However, all other odorants are not tightly clustered across their 3 concentrations, suggesting that the behavioral profile changes with concentration. PCA further supported these results (Figure S4B).

Finally, to compare the magnitude of changes in odorant identity versus changes in odorant concentration, we calculated the Euclidean distances between all possible odorant-concentration pairs. The average distances for different odorant pairs of equal concentrations (i.e., inter-odorant) and for different concentration pairs of the same odorant (i.e., inter-concentration) are not significantly different (Figure 5D). However, the average inter-odorant distances were significantly higher for odorant pairs at the highest concentration (85 mM) compared to pairs at the two lower concentrations (850 μM and 8.5 μM). Odorant identity and concentration thus had a similar quantitative impact on mouse olfactory behavior, with higher odorant concentrations producing greater behavioral diversity.

DISCUSSION

Studies from the last 30 years have contributed to quantification, characterization, and prediction of human olfactory perception. This was enabled by physicochemical descriptions of odorants, novel statistical methods, and machine-learning algorithms applied to large human olfactory psychophysical datasets.⁶⁻¹¹

In comparison, mouse olfactory perception is still poorly understood, mainly due to the lack of a large odor-guided behavioral dataset. Here, we conducted a large-scale study of olfactory-driven mouse behavior by generating the most comprehensive mouse olfactory ethological atlas to date. We scored 18 behaviors in 525 mice across 98 odorant conditions, generating 9,765 data points encompassing 1,764 odorant-behavior interactions.

In humans, odor character is quantified using behavioral methods, such as free labeling, odor profiling, or pairwise similarity.⁴⁸ These methods show that, despite interindividual variation, humans have shared and reproducible olfactory percepts for most odorants. Similarly, we find that mouse olfactory perception can be quantified by measuring several behaviors and that such behaviors demonstrate shared and reproducible olfactory perception across individuals. Many odorants were distinguishable from each other or from the ensemble using such quantified behaviors as observed in a novel mouse. Behavior in novel mice was best reconstructed using a low-dimensional space built from the behavior of other mice. Furthermore, a single dimension showed a modest correlation ($r = 0.5$) between mouse behavior and human perception. Lastly, some mouse behaviors were at least partly predictable from chemical structure alone. Together, these data indicate that there is indeed a rich, canonical set of odor-evoked behaviors in mouse, and this analysis begins to provide insight into mouse olfactory perception.

We found unique behavioral patterns for most odorants and for different concentrations of the same odorant, highlighting the diversity of odor-guided behavioral responses. The olfactory system employs a combinatorial strategy to maximize the discriminability of distinct odorants at different concentrations.^{39,40} This strategy involves hundreds to thousands of OSN/OR subtypes, present in the olfactory mucosa at different abundances.³ Despite this peripheral diversity, it is hypothesized that behavioral and perceptual dimensionality is lower than it first appears.⁶ Our data were consistent with a low-dimensional space for behavior. Approach-avoid behaviors, surrogate for mouse odor valence, were the most important behavioral dimension across our analyses. Analogously, rated pleasantness, a surrogate for odor valence, is the principal perceptual dimension in human odor descriptions.^{7,12,35} Odor valence may thus play a central role in olfactory perception in mammals more generally. However, we offer only a lower bound on odor-evoked behavioral diversity, complexity, and dimensionality; alternative future techniques could capture additional information spanning additional dimensions. Furthermore, olfactory *perception*, resulting from a combinatorial code of peripheral neural activity, could be higher dimensional still. The transformation from perception to behavior may consist of a projection to lower dimension, a computational transformation consistent with olfactory circuitry.⁴⁹

Valence is the principal axis for both mouse behavior and human perception,⁴³ and indeed, olfactory receptor responses in both mice and humans predict human-rated valence.¹² However, this does not imply that specific valence percepts for odorants are shared across species. Indeed, the shared factor that we identified using CCA did not have a strong loading for human-rated pleasantness or any other specific human percept, suggesting that (at least for this molecular panel)

any shared perceptual and behavioral representation of specific odorants across species may have a more multifactorial origin.

In humans, the same odorant can be perceived as pleasant or unpleasant, depending on its concentration. We and others have shown that odorant concentration also impacts mouse olfactory preferences.^{14,25,31,50,51} Here, we extended the publicly available data on this topic significantly (18 behaviors, 12 odorants, and 3 concentrations). We found that behavior within odorant was not fixed across concentrations and that higher concentrations elicited more diverse behavior across odorants. This is consistent with numerically greater OR and glomerular activation by higher odorant concentrations,^{39,41,52} thus increasing potential combinatorial complexity and facilitating downstream pattern separation. Similar to our findings, human studies also show that similarities in odor quality between distinct molecules are inversely related to concentration.⁴⁵ Odor character is not consistent across concentrations and so it is not an intrinsic property of the molecule. Indeed, our analysis suggests that mouse behavior is a consequence of an interaction between molecular structure and concentration.

Although we cannot exclude the possibility that lower concentrations are undetectably weak for some odorants, we observed significant (versus odorless H₂O control) behavioral changes for all odorants except PEA at the lowest concentration (and PEA activates OSNs at concentrations much lower than that).⁵³

Molecular features of odorants are closely linked to psychophysical and behavioral measures of odor valence in human and mouse, respectively.^{7,17,43} Predictive models based on such features can now accurately predict the human olfactory percept of many odorants.^{8,54,55} We employed similar strategies and found that some mouse odor-guided behaviors can also be partly predicted by the physicochemical properties of odorants. The observation that specific physicochemical properties of odorants can predict some behavioral outputs in mice also suggests new possibilities for studying the functional organization of the mouse olfactory system. Subsets of physicochemical properties predictive of odor-guided behaviors could potentially be linked to the spatial organization of ORs/OSN subtypes in the nasal epithelium. Future large-scale experiments focused on connecting the zonal expression patterns for all mouse ORs to the physicochemical descriptors of their respective agonists will be critical to test this hypothesis.

Future studies using recently developed and fully automated techniques, such as MoSeq and DeepLabCut,^{24,56} in combination with behavioral recordings meeting or exceeding the volume and diversity of odorants and mice used here, have the potential to further elucidate the structure of mouse behavioral space and how it relates to odorant identity, valence, and character. We hope that the results presented here will motivate such future work, as we have shown the link between odorant and behavior is strong and shared across individuals.

In conclusion, our study provides a foundational quantitative database of odor-guided behaviors in the mouse that can be exploited in future studies to further deconstruct many aspects of mouse olfactory behavior and putative perception and facilitate future comparative studies of olfactory percepts among different species.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
 - Lead contact
 - Materials availability
 - Data and code availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Animals
- **METHOD DETAILS**
 - Behavioral scoring
 - Machine Learning
 - DREAM predictions
- **QUANTIFICATION AND STATISTICAL ANALYSIS**

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cub.2021.04.020>.

ACKNOWLEDGMENTS

We would like to thank Dr. Darren W. Logan and the members of the Saraiva Lab for the constructive feedback and Alex Williams for the ridge-regularized CCA code. This work was supported by Sidra Medicine (SDR400077), a member of Qatar Foundation, and the National Institute of Health (U19NS112953 and R01DC018455).

AUTHOR CONTRIBUTIONS

D.M. participated in the design of the project, analyzed data, and wrote the initial version of the manuscript. M.M., C.J.A., R.H., A.S., S.D., D.A., and S.S. analyzed data. J.D.M. analyzed data and wrote the final version of the manuscript. R.C.G. designed the data analysis methodology, analyzed data, and wrote the final version of the manuscript. L.R.S. analyzed data, conceived and supervised the project, and wrote the final version of the manuscript.

DECLARATION OF INTERESTS

J.D.M. receives research funding from Google Research and Procter & Gamble and was on the scientific advisory board of Aromyx and received compensation for these activities. R.C.G. receives research funding from Google Research and The Taylor Corporation and is on the advisory board of Climax Foods. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All other authors declare that they have no competing interests.

Received: November 26, 2020

Revised: March 9, 2021

Accepted: April 9, 2021

Published: May 5, 2021

REFERENCES

1. Bear, D.M., Lassance, J.M., Hoekstra, H.E., and Datta, S.R. (2016). The evolving neural and genetic architecture of vertebrate olfaction. *Curr. Biol.* 26, R1039–R1049.
2. Li, Q., and Liberles, S.D. (2015). Aversion and attraction through olfaction. *Curr. Biol.* 25, R120–R129.
3. Saraiva, L.R., Riveros-McKay, F., Mezzavilla, M., Abou-Moussa, E.H., Arayata, C.J., Makhoul, M., Trimmer, C., Ibarra-Soria, X., Khan, M., Van Gerven, L., et al. (2019). A transcriptomic atlas of mammalian olfactory mucosae reveals an evolutionary influence on food odor detection in humans. *Sci. Adv.* 5, eaax0396.
4. Saraiva, L.R., Ahuja, G., Ivandic, I., Syed, A.S., Marioni, J.C., Korsching, S.I., and Logan, D.W. (2015). Molecular and neuronal homology between the olfactory systems of zebrafish and mouse. *Sci. Rep.* 5, 11487.
5. Niimura, Y. (2012). Olfactory receptor multigene family in vertebrates: from the viewpoint of evolutionary genomics. *Curr. Genomics* 13, 103–114.
6. Secundo, L., Snitz, K., and Sobel, N. (2014). The perceptual logic of smell. *Curr. Opin. Neurobiol.* 25, 107–115.
7. Yeshurun, Y., and Sobel, N. (2010). An odor is not worth a thousand words: from multidimensional odors to unidimensional odor objects. *Annu. Rev. Psychol.* 61, 219–241, C1–C5.
8. Keller, A., Gerkin, R.C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., Mainland, J.D., Ihara, Y., Yu, C.W., Wolfinger, R., et al.; DREAM Olfaction Prediction Consortium (2017). Predicting human olfactory perception from chemical features of odor molecules. *Science* 355, 820–826.
9. Keller, A., and Vosshall, L.B. (2016). Olfactory perception of chemically diverse molecules. *BMC Neurosci.* 17, 55.
10. Dravnieks, A. (1982). Odor quality: semantically generated multidimensional profiles are stable. *Science* 218, 799–801.
11. Moskowitz, H.R., Dravnieks, A., and Klarman, L.A. (1976). Odor intensity and pleasantness for a diverse set of odorants. *Percept. Psychophys.* 19, 122–128.
12. Haddad, R., Weiss, T., Khan, R., Nadler, B., Mandaïron, N., Bensafi, M., Schneidman, E., and Sobel, N. (2010). Global features of neural activity in the olfactory system form a parallel code that predicts olfactory behavior and perception. *J. Neurosci.* 30, 9017–9026.
13. Gross-Isseroff, R., and Lancet, D. (1988). Concentration-dependent changes of perceived odor quality. *Chem. Senses* 13, 191–204.
14. Kobayakawa, K., Kobayakawa, R., Matsumoto, H., Oka, Y., Imai, T., Ikawa, M., Okabe, M., Ikeda, T., Itoharu, S., Kikusui, T., et al. (2007). Innate versus learned odour processing in the mouse olfactory bulb. *Nature* 450, 503–508.
15. Root, C.M., Denny, C.A., Hen, R., and Axel, R. (2014). The participation of cortical amygdala in innate, odour-driven behaviour. *Nature* 515, 269–273.
16. Logan, D.W., Brunet, L.J., Webb, W.R., Cutforth, T., Ngai, J., and Stowers, L. (2012). Learned recognition of maternal signature odors mediates the first suckling episode in mice. *Curr. Biol.* 22, 1998–2007.
17. Mandaïron, N., Poncelet, J., Bensafi, M., and Didier, A. (2009). Humans and mice express similar olfactory preferences. *PLoS ONE* 4, e4209.
18. Ferrero, D.M., Lemon, J.K., Fluegge, D., Pashkovski, S.L., Korzan, W.J., Datta, S.R., Spehr, M., Fendt, M., and Liberles, S.D. (2011). Detection and avoidance of a carnivore odor by prey. *Proc. Natl. Acad. Sci. USA* 108, 11235–11240.
19. Ueno, H., Shimada, A., Suemitsu, S., Murakami, S., Kitamura, N., Wani, K., Matsumoto, Y., Okamoto, M., and Ishihara, T. (2019). Anti-depressive-like effect of 2-phenylethanol inhalation in mice. *Biomed. Pharmacother.* 111, 1499–1506.
20. Papes, F., Logan, D.W., and Stowers, L. (2010). The vomeronasal organ mediates interspecies defensive behaviors through detection of protein pheromone homologs. *Cell* 141, 692–703.
21. Brechbühl, J., Moine, F., Klaey, M., Nenniger-Tosato, M., Humi, N., Sporkert, F., Giroud, C., and Broillet, M.C. (2013). Mouse alarm pheromone shares structural similarity with predator scents. *Proc. Natl. Acad. Sci. USA* 110, 4762–4767.
22. Kondoh, K., Lu, Z., Ye, X., Olson, D.P., Lowell, B.B., and Buck, L.B. (2016). A specific area of olfactory cortex involved in stress hormone responses to predator odors. *Nature* 532, 103–106.
23. Saito, H., Nishizumi, H., Suzuki, S., Matsumoto, H., Ieki, N., Abe, T., Kiyonari, H., Morita, M., Yokota, H., Hirayama, N., et al. (2017). Immobility responses are induced by photoactivation of single glomerular species responsive to fox odour TMT. *Nat. Commun.* 8, 16011.

24. Wiltschko, A.B., Johnson, M.J., Iurilli, G., Peterson, R.E., Katon, J.M., Pashkovski, S.L., Abaira, V.E., Adams, R.P., and Datta, S.R. (2015). Mapping sub-second structure in mouse behavior. *Neuron* 88, 1121–1135.
25. Saraiva, L.R., Kondoh, K., Ye, X., Yoon, K.H., Hernandez, M., and Buck, L.B. (2016). Combinatorial effects of odorants on mouse behavior. *Proc. Natl. Acad. Sci. USA* 113, E3300–E3306.
26. Fortes-Marco, L., Lanuza, E., and Martinez-Garcia, F. (2013). Of pheromones and kairomones: what receptors mediate innate emotional responses? *Anat. Rec. (Hoboken)* 296, 1346–1363.
27. Novotny, M.V. (2003). Pheromones, binding proteins and receptor responses in rodents. *Biochem. Soc. Trans.* 31, 117–122.
28. Liberles, S.D. (2014). Mammalian pheromones. *Annu. Rev. Physiol.* 76, 151–175.
29. Achiraman, S., and Archunan, G. (2002). Characterization of urinary volatiles in Swiss male mice (*Mus musculus*): bioassay of identified compounds. *J. Biosci.* 27, 679–686.
30. Apfelbach, R., Soini, H.A., Vasilieva, N.Y., and Novotny, M.V. (2015). Behavioral responses of predator-naïve dwarf hamsters (*Phodopus campbelli*) to odor cues of the European ferret fed with different prey species. *Physiol. Behav.* 146, 57–66.
31. Li, Q., Korzan, W.J., Ferrero, D.M., Chang, R.B., Roy, D.S., Buchi, M., Lemon, J.K., Kaur, A.W., Stowers, L., Fendt, M., and Liberles, S.D. (2013). Synchronous evolution of an odor biosynthesis pathway and behavioral response. *Curr. Biol.* 23, 11–20.
32. Füzesi, T., Daviu, N., Wamsteeker Cusulin, J.I., Bonin, R.P., and Bains, J.S. (2016). Hypothalamic CRH neurons orchestrate complex behaviours after stress. *Nat. Commun.* 7, 11937.
33. Pérez-Gómez, A., Bleymehl, K., Stein, B., Pyrski, M., Birnbaumer, L., Munger, S.D., Leinders-Zufall, T., Zufall, F., and Chamero, P. (2015). Innate predator odor aversion driven by parallel olfactory subsystems that converge in the ventromedial hypothalamus. *Curr. Biol.* 25, 1340–1346.
34. Cichy, A., Shah, A., Dewan, A., Kaye, S., and Bozza, T. (2019). Genetic depletion of class I odorant receptors impacts perception of carboxylic acids. *Curr. Biol.* 29, 2687–2697.e4.
35. Haddad, R., Carmel, L., Sobel, N., and Harel, D. (2008). Predicting the receptive range of olfactory receptors. *PLoS Comput. Biol.* 4, e18.
36. Lee, D.D., and Seung, H.S. (1999). Learning the parts of objects by non-negative matrix factorization. *Nature* 401, 788–791.
37. Castro, J.B., Ramanathan, A., and Chennubhotla, C.S. (2013). Categorical dimensions of human odor descriptor space revealed by non-negative matrix factorization. *PLoS ONE* 8, e73289.
38. Manoel, D., Makhoul, M., Scialdone, A., and Saraiva, L.R. (2019). Interspecific variation of olfactory preferences in flies, mice, and humans. *Chem. Senses* 44, 7–9.
39. Malnic, B., Hirono, J., Sato, T., and Buck, L.B. (1999). Combinatorial receptor codes for odors. *Cell* 96, 713–723.
40. Nara, K., Saraiva, L.R., Ye, X., and Buck, L.B. (2011). A large-scale analysis of odor coding in the olfactory epithelium. *J. Neurosci.* 31, 9179–9191.
41. Johnson, B.A., and Leon, M. (2000). Modular representations of odorants in the glomerular layer of the rat olfactory bulb and the effects of stimulus concentration. *J. Comp. Neurol.* 422, 496–509.
42. Haddad, R., Khan, R., Takahashi, Y.K., Mori, K., Harel, D., and Sobel, N. (2008). A metric for odorant comparison. *Nat. Methods* 5, 425–429.
43. Khan, R.M., Luk, C.H., Flinker, A., Aggarwal, A., Lapid, H., Haddad, R., and Sobel, N. (2007). Predicting odor pleasantness from odorant structure: pleasantness as a reflection of the physical world. *J. Neurosci.* 27, 10015–10023.
44. Licon, C.C., Bosc, G., Sabri, M., Mantel, M., Fournel, A., Bushdid, C., Golebiowski, J., Robardet, C., Plantevit, M., Kaytue, M., and Bensafi, M. (2019). Chemical features mining provides new descriptive structure-odor relationships. *PLoS Comput. Biol.* 15, e1006945.
45. Laing, D.G., Legha, P.K., Jinks, A.L., and Hutchinson, I. (2003). Relationship between molecular structure, concentration and odor qualities of oxygenated aliphatic molecules. *Chem. Senses* 28, 57–69.
46. Sirotnin, Y.B., Shusterman, R., and Rinberg, D. (2015). Neural coding of perceived odor intensity. *eNeuro* 2, ENEURO.0083-15.2015.
47. Jain, A.K., and Moreau, J.V. (1987). Bootstrap technique in cluster analysis. *Pattern Recognit.* 20, 547–568.
48. Wise, P.M., Olsson, M.J., and Cain, W.S. (2000). Quantification of odor quality. *Chem. Senses* 25, 429–443.
49. Pashkovski, S.L., Iurilli, G., Brann, D., Chicharro, D., Drummey, K., Franks, K.M., Panzeri, S., and Datta, S.R. (2020). Structure and flexibility in cortical representations of odour space. *Nature* 583, 253–258.
50. Sato-Akuhara, N., Horio, N., Kato-Namba, A., Yoshikawa, K., Niimura, Y., Ihara, S., Shirasu, M., and Touhara, K. (2016). Ligand specificity and evolution of mammalian musk odor receptors: effect of single receptor deletion on odor detection. *J. Neurosci.* 36, 4482–4491.
51. Horio, N., Murata, K., Yoshikawa, K., Yoshihara, Y., and Touhara, K. (2019). Contribution of individual olfactory receptors to odor-induced attractive or aversive behavior in mice. *Nat. Commun.* 10, 209.
52. Fried, H.U., Fuss, S.H., and Korsching, S.I. (2002). Selective imaging of presynaptic activity in the mouse olfactory bulb shows concentration and structure dependence of odor responses in identified glomeruli. *Proc. Natl. Acad. Sci. USA* 99, 3222–3227.
53. Zhang, J., Pacifico, R., Cawley, D., Feinstein, P., and Bozza, T. (2013). Ultrasensitive detection of amines by a trace amine-associated receptor. *J. Neurosci.* 33, 3228–3239.
54. Kowalewski, J., Huynh, B., and Ray, A. (2021). A systems-wide understanding of the human olfactory percept chemical space. *Chem. Senses*. Published online February 26, 2021. <https://doi.org/10.1093/chemse/bjab007>.
55. Sanchez-Lengeling, B., Wei, J.N., Lee, B.K., Gerkin, R.C., Aspuru-Guzik, A., and Wiltschko, A.B. (2019). Machine learning for scent: learning generalizable perceptual representations of small molecules. *arXiv*, arXiv:1910.10685v2. <https://arxiv.org/abs/1910.10685>.
56. Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat. Neurosci.* 21, 1281–1289.
57. Benjamini, Y., Krieger, A.M., and Yekutieli, D. (2006). Adaptive linear step-up procedures that control the false discovery rate. *Biometrika* 93, 491–507.
58. Williams, A. (2020). Ridge CCA. <https://gist.github.com/ahwillia/f7d6d27e176df65d6baa6aee8b661042>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental models: Organisms/strains		
Mouse: C57BL/6J	The Jackson Laboratory	JAX: 000664
Software and algorithms		
Code generated here	This manuscript	https://github.com/rgerkin/manoel-2021
Ethovision XT	Noldus	https://www.noldus.com/ethovision-xt
Dragon 6.0	Talete	http://www.talete.mi.it/
Python 3	Python	http://python.org
Pandas	AQR Capital Management	https://pandas.pydata.org/
Scikit-Learn	scikit-learn	https://scikit-learn.org/stable/
Pyrfume	The Pyrfume Project	https://pyrfume.org
Mordred	Mordred Descriptors	https://github.com/ordred-descriptor/mordred
Rdkit	Rdkit	http://www.rdkit.org
R Statistical Software 3.5.1	R project for Statistical Computing	https://www.r-project.org/
GraphPad Prism	Graphpad	https://www.graphpad.com/scientific-software/prism/
Paleontological Statistics	Paleontological Museum, University of Oslo	https://folk.universitetetioslo.no/ohammer/past
Other		
Behavior video library	Saraiva et al. ²⁵	N/A

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and data should be directed to and will be fulfilled by the Lead Contacts, Luis R. Saraiva (saraivalmr@gmail.com) or Richard C. Gerkin (email: rgerkin@asu.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

The published article includes all datasets generated or analyzed during this study, and the code generated during this study is available at <https://github.com/rgerkin/manoel-2021>.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

The animals used in the video library from our previous study²⁵ were adult male C57BL/6J mice (aged 8–14 weeks, The Jackson Laboratory). Each mouse was randomly assigned and exposed to only a single odorant, and thus the data for each odorant consisted of 5–10 mice each scored on 18 behaviors. The experiments performed in Saraiva et al.²⁵ were approved by the Fred Hutchinson Cancer Research Center Institutional Animal Care and Use Committee.

METHOD DETAILS

Behavioral scoring

We retrieved a video library from our previous study,²⁵ in which we subjected adult male mice to the olfactory preference test for a total duration of 3 minutes. In this study we analyzed 410 videos from mice exposed to an odorless control (i.e., water, or H₂O) or one of the 73 odorants at a single concentration (at 85 mM), and 123 additional videos from mice exposed to two other descending concentrations (850 μ M and 8.5 μ M) for a subset of 12 odorants. The odorants tested in the videos include 61 general odorants and 12

ethologically relevant odorants (7 mouse pheromones, and 5 kairomones). All these compounds were chemically diverse and belong to multiple chemical structural classes (underlined, for more details see Saraiva et al.²⁵) as shown below, followed by their 3-letter abbreviation in parentheses: Alcohols: 2-phenylethanol (2PE); geraniol (GER); heptanol (HPO); hexanol (HXO); linalool (LIN); octanol (OCO); cis-3-hexenol (C3H). Aldehydes: benzaldehyde (BZL); citral (CIT); citronellal (CTN); heptanal (HPN); octanal (OCN); trans-2-hexenal (T2H). Amines: 2-methylbutylamine (2MB); 1-(2-aminoethyl)piperidine (AEP); aniline (ANI); 3-amino-s-triazole (AST); cadaverine (CAD); N,N-dimethylbutylamine (DMB); N,N-dimethylethylamine (DME); N,N-dimethyloctylamine (DMO); N,N-dimethylcyclohexylamine (DMC); heptylamine (HEP); hexylamine (HXA); isoamylamine (IAA); 2-methyl-1-pyrroline (M1P); N-methylpiperidine (NMP); octylamine (OCT); 2-phenylethylamine (PEA); trimethylamine (TMA); putrescine (PUT); pyrrolidine (PYR); spermidine (SPD); spermine (SPN); o-toluidine (TOL). Azines: 2,5-dimethylpirazine (DMP); 2-ethyl-3,5(6)-dimethylpirazine (EDM); indole (IND); quinoline (QUI); skatole (SKA). Camphors: (+/-)-camphor (CAM); (-)-fenchone (-FCH); (+)-fenchone (+FCH); eucalyptol (EUC). Carboxylic acids: 2-methylbutyric acid (MBA); octanoic acid (OCA); propionic acid (PPA). Esters: amyl acetate (AAC); ethyl butyrate (EBT). Ketones: 2-heptanone (2HO); alpha-ionone (ION). Musks: ambrettolide (AMB); civettone (CIV); muscone (MUS). Terpenes: beta-farnesene (BFA); (-)-carvone (-CVN); (+)-carvone (+CVN); farnesene mixed isomers (alpha+beta) (FAR); (-)-limonene (-LIM); (+)-limonene (+LIM); (+)-menthol (+MEN); (-)-menthone (-MNT); (+)-menthone (+MNT); alpha-pinene (PIN); rose oxide (ROX). Thiazoles: 2-isobutylthiazole (IBT); 2-isopropyl-4-5-dihydrothiazole (IPT); 2,5-dihydro-2,4,5-trimethylthiazoline (TMT). Thiols: 2-propylthietane (2PT); hexanethiol (HXT); octanethiol (OTT). Vanillin-like compounds: eugenol (EUG); vanillin (VAN).

For each video, in addition to the cumulative duration of olfactory investigation (3'dOI) reported in Saraiva et al.²⁵ for the 3 minutes-long assay, we scored 17 new behavioral parameters indicative of either valence, stress, and exploration. The videos were randomized and scored blind (to the odorant) using the following criteria:

Olfactory investigation (OI) parameters

1'dOI, 1'fOI, and 3'fOI represent the cumulative duration (d) or frequency (f) of olfactory investigation during the 1st minute (1') or the full 3 minutes (3') of the assay. We considered OI only if the nose of the mouse was overlapping, or in very close proximity (~0.5 cm) of the stimulus. For these parameters, the videos were randomized and scored by the experimenter.

Zone (Z) parameters

1'Z1, 1'Z3, 3'Z1, and 3'Z3 represent the cumulative time mice spent in either Zone 1 (Z1) or Zone 3 (Z3) of the cage during the 1st minute (1') or the full 3 minutes (3') of the assay. Here, the test cage was divided into three equal-sized zones, with Z1 representing the zone furthest away from the odor stimulus and Z3 the zone containing the stimulus. Time spent inside each zone (head of the animal had to be within the zone) was scored using the Ethovision XT software (version 11, Noldus Information Technology), and videos in which the mouse transported the piece of filter paper outside Z3 were not included. For these parameters, the videos were scored by a non-experimenter.

Handling (H) or catching (C) parameters

dH/C and fH/C represent the cumulative duration (d) or frequency (f) where mice handled and/or caught the stimulus with its front paws during the 3 minutes of the assay. For these parameters, the videos were scored by a non-experimenter.

Risk assessment (RAS)

This parameter represents the number of episodes the mouse displays the flat-back/stretch-attend response, followed by a sniff in the direction of the stimulus, during the 3 minutes of the assay. For this parameter, the videos were scored by a non-experimenter.

Escape (ESC)

This parameter represents the number of episodes the mouse displays a quick and contactless approach toward the stimulus, followed by an even faster withdrawal/darting to the opposite end of the cage, during the 3 minutes of the assay. For this parameter, the videos were scored by a non-experimenter.

Digging (DIG) parameters

dDIG and fDIG represent the cumulative duration (d) or frequency (f) that the mouse digs into the bedding with the forelimbs, often kicking it away with the hindlimbs, during the 3 minutes of the assay. For these parameters, the videos were scored by a non-experimenter.

Distance (DIT)

This parameter represents the total length the mouse walked/ran through during the three minutes duration of the video. Distance traveled was scored using the Ethovision XT software (version 11, Noldus Information Technology), and video tracking done using the center-point of the mouse. For this parameter, the videos were scored by a non-experimenter.

Rearing (REA)

This parameter represents the number of episodes the mouse stands on its hindlegs (rearing) anywhere in the cage, including when rearing against the walls during the 3 minutes of the assay. For this parameter, the videos were scored by a non-experimenter.

Self-grooming (SGR) parameters

dSGR and fSGR represent the cumulative duration (d) or frequency (f) where the mice are self-grooming, defined by when the mouse was licking its fur, grooming itself with the forepaws, or scratching any part of its body with any limb. For these parameters, the videos were scored by a non-experimenter.

Machine Learning

We used Python 3.8 and the Pyrfume package (version 0.15) to calculate 1826 physicochemical descriptors for 74 molecules (Mordred, version 1.2). Descriptors with zero variance or > 25% missing values were removed leaving 1328 descriptors, and the remaining

missing values were iteratively imputed using (scikit-learn, version 0.23.2). We also used Pyrfume to calculate the Morgan fingerprint similarity (rdkit, version 2020.09.1) of the 74 molecules to a reference set of 9645 (mostly) odorous compounds. We used support vector regression with leave-one-out cross-validation (scikit-learn) to train two models, one based on Mordred descriptors and one on Morgan similarities, and their predictions were averaged before validation. We evaluated two kinds of predictions: (1) One where the models were trained on all but a single mouse, i.e., the training set included other instances of the same odorant presented to different mice, and (2) one where the models were trained on all but a single odorant, i.e., the training set excluded all mice presented with that odorant. For (1) evaluation was done per mouse and for (2) averaging across mice within odorant was performed first and training/testing was done on these averages. We reported the Pearson correlation r between the observed (out-of-sample) behavior and the predicted behavior; r was chosen to facilitate direct comparison to several predictive models in human olfactory perception that also use this measure.^{8,54,55} Significance was assessed by converting r values to Z-scores using the Fisher transformation and using the standard error for the Fisher Z of $1/\sqrt{n-3}$ in conjunction with the cumulative normal distribution. We did not correct our reported values for multiple comparisons, but a False Discovery Rate correction would have had minimal impact.

DREAM predictions

The DREAM model developed in Keller et al.⁸ was applied to H₂O and each of the 73 odorants used in the current study to generate predictions for each of 21 perceptual descriptors. Briefly, an isomeric, canonical SMILES string was generated for each odorant using rdkit (Python) and used to generate a rdkit mol object in which 3-dimensional coordinates of each atom position were estimated. In order to match the methods of Keller et al.,⁸ Dragon 6.0 was used to compute features from these 3-dimensional structures. The DREAM model was re-trained from scratch on the 476 original molecules (from Keller et al.⁸) and used to predict 21 perceptual descriptors from the 74 odorants used in the current study.

QUANTIFICATION AND STATISTICAL ANALYSIS

The number of mice tested for each odorant and the quantification of each behavioral measurement are described in the “Behavioral Scoring” section above. The raw behavioral scores and numbers of animals used can be consulted in [Data S1](#). One-way ANOVA tests were performed using GraphPad Prism 8.0.0 software for each of the 18 behavioral parameters. P values for the pairwise comparisons between H₂O control and each of the tested odorant stimuli were computed with a two-stage Benjamini, Krieger and Yekutieli (BKY) multiple comparisons correction⁵⁷ ([Figures 1E and 5A](#); [Data S1](#)).

Correlation matrices generated with pandas and scikit-learn were used to compute the eigenvalues and eigenvectors in the principal components analysis ([Figures 1F and 1G](#)). For [Figure S1C](#), the odorant coloring was done by computing the first three principal components of the Morgan fingerprint similarity matrix (computed using rdkit and pyrfume), projecting the odorants onto this 3d-space, normalizing each dimension between 0 and 1, and setting the⁵⁴ values of each odorant according to these normalized values.

Custom code using numpy and scipy was used to compute multivariate D' ([Figures 2A–2D](#)) and scikit-learn was used to train the linear classifiers ([Figures 2E–2I](#)). Scikit-learn was used to perform the NMF analysis ([Figures 3A and 3B](#)) and custom code⁵⁸ was used to implement the ridge-regularized CCA ([Figure 3C](#)).

Spearman's correlation coefficients of the 1,536 physicochemical descriptors against the 18 behavioral parameters were computed using Graphpad Prism 8.0.0 software ([Figure 4C](#)).

The standardized data (z-scores) in [Figure S4A](#) were further analyzed through hierarchical clustering performed with PALaeontological STatistics (version 4.06) using Euclidean distances with Ward's method ([Figure 5C](#)). The z-score data matrix was also used for principal component analyses and performed using PALaeontological STatistics (version 4.06) ([Figure S4B](#)).

Current Biology, Volume 31

Supplemental Information

Deconstructing the mouse olfactory percept through an ethological atlas

Diogo Manoel, Melanie Makhlouf, Charles J. Arayata, Abbirami Sathappan, Sahar Da'as, Doua Abdelrahman, Senthil Selvaraj, Reem Hasnah, Joel D. Mainland, Richard C. Gerkin, and Luis R. Saraiva




Figure S1. An olfactory-ethological atlas and primary axis of olfactory perception in the mouse, related to Figure 1. (A) Heatmap summarizing the 7380-individual z-scores representing of 18 mouse behavioral responses to 73 odorants ($n = 410$ mice). **(B)** Cumulative variance explained by the principal components (PCs). **(C)** Principal Component Analysis of the 18 mouse behavioral parameters for H₂O and the 73 tested odorants ($n = 3$ -7 per odorant). Circles are colored according to their chemical structure similarity, which was computed using Morgan fingerprints (see STAR Methods). See also Figure 1, and Data S1.




Figure S2. Details of the non-negative matrix factorization (NMF) used in Figure 3, related to Figures 2 and 3. (A) Behavioral loadings for each of the first two NMF factors. Each point reflects the loading for one of the 18 behaviors. **(B)** Projection of the behavioral data for all odorants into a space comprising these factors. Each point reflects one of the 73 odorants. These two factors are highly negatively correlated, suggesting that they mostly describe opposing valence factors (e.g. approach vs avoid). See also Figures 2, S3, and Table S1.




Figure S3. Comparing the mouse and human olfactory percepts, related to Figures 3 and 4. (A) Correlogram between the 18 mouse behavioral parameters and 21 human ratings for the 23 odorants overlapping between this study and ^{S1}. Only significant ($P < 0.05$) correlations are plotted, and the circle size and color indicate the magnitude and direction of the correlation (Spearman ρ). Blank cells correspond to non-significant correlations. **(B)** Correlation between each of the 21 human-rated perceptual descriptors for 73 odorants (the 22 odorants common to this study and ^{S1} as well as the 51 odorants predicted using ^{S2} and i) the descriptors themselves and ii) the principal components from Fig. 1. **(C)** Same as Fig S3B, but for NMF factors in Fig. 3. See also Figures 2, 4, S2, and Table S1.




Figure S4. The effect of odorant dosage on behavior, related to Figure 5. (A) Heatmap indicating the z-score for each of 12 odorants (relative to all other odorants) for the 18 behavioral parameters measured after exposure to those odorants at three concentrations (85mM, 850μM, and 8.5μM, see STAR Methods). **(B)** Principal component analysis of the behavioral profiles for the 12 odorants tested at the three different concentrations. H₂O is shown in blue, aversive odorants in magenta, neutral in grey, and approached in green. The line connecting the circles indicates the minimum spanning tree. See also Figure 5, and Data S3.

Human parameters	Mean	StDev	Z
Acid	0.048702194	0.07531571	0.646640585
Ammonia	-0.009887358	0.066986152	-0.147603016
Bakery	-0.037669011	0.060638897	-0.621202118
Burnt	-0.121437285	0.076369271	-1.590132828
Chemical	0.039923889	0.071582565	0.557732029
Cold	-0.011707885	0.070135746	-0.166931781
Decayed	-0.023797411	0.081189952	-0.293107834
Fish	0.005314301	0.087729115	0.060576247
Flower	-0.07325992	0.06768923	-1.082298035
Fruit	0.023629055	0.066334428	0.356211031
Garlic	0.102715311	0.085814371	1.196947672
Grass	0.011990264	0.054774955	0.218900474
Intensity	0.058308333	0.059595577	0.97840034
Musky	-0.005165428	0.0783955	-0.065889347
Pleasantness	-0.081756002	0.09275403	-0.881428029
Sour	-0.081562531	0.078791444	-1.035169908
Spices	-0.078923349	0.068496048	-1.152232154
Sweaty	-0.01983751	0.092195631	-0.215167576
Sweet	0.039596144	0.080172095	0.493889353
Warm	0.065088094	0.085555097	0.760774008
Wood	0.054981654	0.060035691	0.915816124

Table S1. Weights of each human perceptual descriptor in the first CCA dimension aligning human perceptual and mouse behavioral spaces, related to Figure 3. The standard deviation is obtained using 100 shuffles of the original data, applying CCA to each, and computing the standard deviation across weights obtained from these shuffles. See also Figures 3, S2, and S3.

SUPPLEMENTAL REFERENCES

S1. Keller, A., and Vosshall, L.B. (2016). Olfactory perception of chemically diverse molecules. *BMC neuroscience* 17, 55.

S2. Keller, A., Gerkin, R.C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., Mainland, J.D., Ihara, Y., Yu, C.W., Wolfinger, R., et al. (2017). Predicting human olfactory perception from chemical features of odor molecules. *Science* 355, 820-826.